arXiv:1407.5745v1 [math.GN] 22 Jul 2014

DIAGONALS OF SEPARATELY CONTINUOUS FUNCTIONS AND THEIR ANALOGS

OLENA KARLOVA, VOLODYMYR MYKHAYLYUK, AND OLEKSANDR SOBCHUK

Abstract. We prove that for a topological space X, an equiconnected space Z and a Baire-one mapping $g : X \to Z$ there exists a separately continuous mapping $f : X^2 \to Z$ with the diagonal g, i.e. $g(x) = f(x, x)$ for every $x \in X$. Under a mild assumptions on X and Z we obtain that diagonals of separately continuous mappings $f : X^2 \to Z$ are exactly Baire-one functions, and diagonals of mappings $f : X^2 \to Z$ which are continuous on the first variable and Lipschitz (differentiable) on the second one, are exactly the functions of stable first Baire class.

1. Introduction

Let $f : X^2 \to Y$ be a mapping. We call a mapping $g : X \to Y$, $g(x) = f(x, x)$, *the diagonal of* f.

Investigations of diagonals of separately continuous functions $f : X^2 \to \mathbb{R}$ started in the classical work of R. Baire [1], who showed that diagonals of separately continuous functions of two real variables are exactly Baire-one functions, i.e. pointwise limits of continuous functions. His result was generalized by A. Lebesgue and H. Hahn for real-valued functions of n real variables (see [9, 10, 6]).

Since the second half the XX-th century, Baire classification of separately continuous mappings and their analogs was intensively studied by many mathematicians (see [13, 15, 16, 12, 2, 5]). Let us observe that W. Rudin [15] was the first who applied partitions of unity to classify separately continuous functions defined on the product of a metrizable space and a topological space and with values in locally convex spaces. Later this method has been developed and now a partition of unity is a familiar tool for Baire classification of separately continuous functions.

A problem on construction of separately continuous functions of n variables with the given diagonal of $(n-1)$-th Baire class was solved in [11]. In [14] it was shown that for any topological space X and a function $g : X \to \mathbb{R}$ of $(n-1)$-th Baire class there exist a separately continuous function $f : X^n \to \mathbb{R}$ with the diagonal g.

Also naturally appear problems on diagonals of CL- and CD-mappings, i.e. diagonals of mappings $f : X^2 \to Y$ which are continuous with respect to the first variable and Lipschitz or differentiable with respect to the second variable.

In the given paper we generalize the result from [14] which concerns a construction of separately continuous mappings $f : X^2 \to Z$ with a given diagonal of the first Baire class to the case of an equiconnected space Z. We also describe diagonals of CL- and CD-mappings for wide classes of spaces X and Z.

2. Preliminaries

By $P(X, Y)$ we will denote the collection of all mappings $f : X \to Y$ with a property P. In particular, by $C(X, Y)$ we denote the collection of all continuous mappings between topological spaces X and Y.

For a mapping $f : X \times Y \to Z$ and a point $(x, y) \in X \times Y$ we write

$$f^x(y) = f_y(x) = f(x, y).$$

Let P and Q be some properties of mappings. We will use the symbol $PQ(X \times Y, Z)$ to denote the collection of all mappings $f : X \times Y \to Z$ which have a property P with respect to the first variable and a property Q with respect to the second one, i.e. $\{y \in Y : f_y \in P(X, Z)\} = Y$ and $\{x \in X : f^x \in Q(Y, Z)\} = X$. If $f \in PP(X \times Y, Z)$, then f is called *a mapping which separately has a property* P. So, for topological spaces X, Y and Z mappings from the class $CC(X \times Y, Z)$ are said to be *separately continuous*.

Recall that a mapping f between topological spaces X and Y is called *a mapping of the first Baire class* or *a Baire-one mapping* if there exists a sequence $(f_n)_{n=1}^{\infty}$ of continuous mappings $f_n : X \to Y$ which is pointwise convergent to f, i.e. $\lim\limits_{n\to\infty} f_n(x) = f(x)$ for every $x \in X$. The collection of all Baire-one mappings $f : X \to Y$ is denoted by $B_1(X, Y)$.

An equiconnected space is a pair (X, λ), consisting of a topological space X and a continuous function $\lambda : X \times X \times [0, 1] \to X$ such that

(i) $\lambda(x, y, 0) = x$;

(ii) $\lambda(x, y, 1) = y$;

(iii) $\lambda(x,x,t)=x$

for all $x,y\in X$ and $t\in[0,1]$.

The simplest example of an equiconnected space is a convex subset of a topological vector space with λ defined by $\lambda(x,y,t)=(1-t)x+ty$.

For a metric space X by $|\cdot-\cdot|_X$ we denote the metric on this space.

3. A method of constructing of separately continuous mappings

In the following theorem we describe a method of a construction of separately continuous mappings with a given diagonal.

Theorem 3.1. *Let X be a topological space, (Z,λ) a metric equiconnected space, let $(G_n)_{n=0}^{\infty}$ and $(F_n)_{n=0}^{\infty}$ be sequences of open sets G_n and closed sets F_n in X^2, $(\varphi_n:X^2\to[0,1])_{n=1}^{\infty}$ be a sequence of separately continuous functions, $(g_n:X\to Z)_{n=1}^{\infty}$ be a sequence of continuous functions pointwise convergent to a function $g:X\to Z$, and $(\delta_n:X\to(0,+\infty))_{n=1}^{\infty}$ be a sequence of positive functions pointwise convergent to zero such that:*

(1) *$G_0=F_0=X^2$ and $\Delta=\{(x,x):x\in X\}\subseteq G_{n+1}\subseteq F_n\subseteq G_n$ for every $n\in\mathbb{N}$;*

(2) *$X^2\setminus G_n\subseteq\varphi_n^{-1}(0)$ and $F_n\subseteq\varphi_n^{-1}(1)$ for every $n\in\mathbb{N}$;*

(3) *$|g_n(x)-g_n(y)|_Z\le\delta_n(y)$ and $|g_{n+1}(x)-g_{n+1}(y)|_Z\le\delta_n(y)$ for every $n\in\mathbb{N}$ and $(x,y)\in G_n$.*

Then the mapping $f:X^2\to Z$,

$$f(x,y)=\begin{cases}\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y)), & (x,y)\in F_{n-1}\setminus F_n\\ g(x), & (x,y)\in E=\bigcap\limits_{n=1}^{\infty}G_n.\end{cases}\tag{3.1}$$

is separately continuous.

Proof. We fix $n\in\mathbb{N}$ and show that

$$f(x,y)=\lambda(\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y)),g_{n+2}(x),\varphi_{n+1}(x,y))\tag{3.2}$$

for all $(x,y)\in F_{n-1}\setminus F_{n+1}$.

Let $(x,y)\in F_{n-1}\setminus F_n$. Since $G_{n+1}\subseteq F_n$, $(x,y)\notin G_{n+1}$ and (2) imply that $\varphi_{n+1}(x,y)=0$. Therefore,

$$\lambda(\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y)),g_{n+2}(x),\varphi_{n+1}(x,y))=$$
$$=\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y))=f(x,y).$$

Now let $(x,y)\in F_n\setminus F_{n+1}$. According to (2), $\varphi_n(x,y)=1$ and

$$\lambda(\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y)),g_{n+2}(x),\varphi_{n+1}(x,y))=$$
$$=\lambda(g_{n+1}(x),g_{n+2}(x),\varphi_{n+1}(x,y))=f(x,y).$$

Since the mappings λ, g_n, g_{n+1}, g_{n+2} are continuous, φ_n and φ_{n+1} are separately continuous, f is continuous with respect to every variable (separately continuous) at each point of the open set $G_n\setminus F_{n+1}$ for every $n\in\mathbb{N}$ as a composition of continuous mappings. Moreover, f is separately continuous at every point of the open set $G_0\setminus F_1=F_0\setminus F_1$. Therefore, f is separately continuous on the open set $X^2\setminus E=\bigcup\limits_{n=1}^{\infty}(G_{n-1}\setminus F_n)$.

It remains to check the continuity of f with respect to x and y at the points of E. We first notice that $(x,y)\in G_n$ for any point $(x,y)\in E$, hence, in accordance with (4), $|g_n(x)-g_n(y)|_Z\le\delta_n(y)$ for every $n\in\mathbb{N}$. After letting $n\to\infty$, we conclude $g(y)=g(x)$.

Fix $(x_0,y_0)\in E$, $\varepsilon>0$ and denote $z_0=f(x_0,y_0)=g(x_0)=g(y_0)$. Since λ is continuous and $\lambda(z_0,z_0,\alpha)=z_0$ for every $\alpha\in[0,1]$, there exists a neighborhood $W=\{z\in Z:|z-z_0|_Z<\delta\}$ of z_0 such that $|\lambda(z_1,z_2,\alpha)-z_0|_Z<\varepsilon$ for any $z_1,z_2\in W$ and $\alpha\in[0,1]$. Since $\lim\limits_{n\to\infty}g_n(x_0)=\lim\limits_{n\to\infty}g_n(y_0)=z_0$ and $\lim\limits_{n\to\infty}\delta_n(y_0)=0$, there exists a number $n_0\in\mathbb{N}$ such that $g_n(x_0)\in W$, $|g_n(y_0)-z_0|_Z<\frac{\delta}{2}$ and $\delta_n(y_0)<\frac{\delta}{2}$ for every $n\ge n_0$. Then we choose neighborhoods U_0 and V_0 of the points x_0 and y_0, respectively, such that $A=(U_0\times\{y_0\})\cup(\{x_0\}\times V_0)\subseteq G_{n_0}$ and show that $|f(x,y)-z_0|_Z<\varepsilon$ for every $(x,y)\in A$.

Let $(x,y)\in A\cap E$. If $y=y_0$, then $f(x,y_0)=g(x)=g(y_0)=z_0$, and if $x=x_0$, then $f(x,y_0)=g(x_0)=z_0$. Hence, $f(x,y)=z_0$ and $|f(x,y)-z_0|_Z<\varepsilon$.

Let $(x,y_0)\in A\setminus E$. We choose a number $n\in\mathbb{N}$ such that $(x,y_0)\in G_n\setminus G_{n+1}$. Since $(x,y_0)\in G_{n_0}$, $n\ge n_0$. According to (3.2),

$$f(x,y_0)=\lambda(g_n(x),g_{n+1}(x),\varphi_n(x,y_0)).$$

Moreover, by (3) and by choosing of n_0, $|g_n(x)-g_n(y_0)|_Z\le\delta_n(y_0)<\frac{\delta}{2}$ and $|g_{n+1}(x)-g_{n+1}(y_0)|_Z\le\delta_n(y_0)<\frac{\delta}{2}$. Taking into account that $|g_n(y_0)-z_0|_Z<\frac{\delta}{2}$ and $|g_{n+1}(y_0)-z_0|_Z<\frac{\delta}{2}$, we deduce that $g_n(x),g_{n+1}(x)\in W$. Therefore, $|f(x,y_0)-z_0|_Z<\varepsilon$.

Now let $(x_0, y) \in A \setminus E$, $n \geq n_0$ and $(x_0, y) \in G_n \setminus G_{n+1}$. Then $f(x_0, y) = \lambda(g_n(x_0), g_{n+1}(x_0), \varphi_n(x_0, y))$, $g_n(x_0), g_{n+1}(x_0) \in W$ and $|f(x_0, y) - z_0|_Z < \varepsilon$.

Hence, f is continuous with respect to the variables x and y at the points of the set E and the proof is complete. □

Recall that a subset A of a topological space X is called *a zero (co-zero) set* if there exists a continuous function $f : X \to [0, 1]$ such that $A = f^{-1}(0)$ ($A = f^{-1}((0, 1])$). Zero and co-zero sets are alternatively called *functionally closed* and *functionally open sets*, respectively.

Corollary 3.2. *Let X be a topological space, (Z, λ) a metrizable equiconnected space and $g \in B_1(X, Z)$. Then there exists a separately continuous mapping $f : X^2 \to Z$ with the diagonal g.*

Proof. Fix any metric $|\cdot - \cdot|_X$ on X which generates its topological structure. Choose a sequence $(g_n)_{n=1}^{\infty}$ of continuous mappings $g_n : X \to Z$ which is pointwise convergent to g. Set $G_0 = F_0 = X^2$,

$$G_n = \{(x, y) \in X^2 : |g_k(x) - g_k(y)|_Z < \frac{1}{n} \quad \forall\, k \leq n + 1\}$$

and

$$F_n = \{(x, y) \in X^2 : |g_k(x) - g_k(y)|_Z \leq \frac{1}{n+1} \quad \forall\, k \leq n + 1\}.$$

Let us observe that the sequences $(g_n)_{n=1}^{\infty}$, $(G_n)_{n=0}^{\infty}$ and $(F_n)_{n=0}^{\infty}$ satisfy conditions (1) and (3) of Theorem 3.1. Moreover, all the sets G_n and F_n are co-zero and zero sets, respectively. Then for every $n \in \mathbb{N}$ there exists a continuous function $\varphi_n : X^2 \to [0, 1]$ with $\varphi_n^{-1}(0) = X^2 \setminus G_n$ and $\varphi_n^{-1}(1) = F_n$. Clearly, the sequence $(\varphi_n)_{n=1}^{\infty}$ satisfies (2). Now Theorem 3.1 implies the existence of a separately continuous mapping f with the diagonal g. □

The following result gives a characterization of diagonals of separately continuous mappings.

Theorem 3.3. *Let X be a topological space, (Z, λ) a metrizable equiconnected space, $g : X \to Z$ a mapping and let, at least one of the following conditions hold:*

(1) *$CC(X \times X, Z) \subseteq B_1(X^2, Z)$;*
(2) *X is a metrically quarter-stratifiable strongly countably-dimensional paracompact space;*
(3) *X is a metrically quarter-stratifiable space and Z is locally convex;*
(4) *X is a metrizable space.*

Then the following conditions are equivalent:

(*i*) *$g \in B_1(X, Z)$;*
(*ii*) *there exists a separately continuous mapping $f : X^2 \to Z$ with the diagonal g.*

Proof. (1). The implication $(i) \Rightarrow (ii)$ follows from Corollary 3.2, and the implication $(ii) \Rightarrow (i)$ follows from the inclusion $CC(X \times X, Z) \subseteq B_1(X^2, Z)$.

(2) and (3). In the case when X is a metrically quarter-stratifiable strongly countably-dimensional paracompact space or X is a metrically quarter-stratifiable space and Z is a locally convex equiconnected space (see definitions in [2]), the inclusion $CC(X \times X, Z) \subseteq B_1(X^2, Z)$ follows from [2, Corollary 5.2].

(4). Notice that an equiconnected space Z is locally arcwise connected. Therefore, by [8], the inclusion $CC(X \times X, Z) \subseteq B_1(X^2, Z)$ holds.

□

The conditions (1) – (4) of Theorem 3.1 suggest the following question.

Question 3.4. *Let X be a quarter-stratifiable space and let Z be a metrizable path-connected space. Does the inclusion $CC(X \times X, Z) \subseteq B_1(X^2, Z)$ hold?*

Let us remark that in the previous theorem the Baire classification of diagonals follows from the stronger inclusion $CC \subseteq B_1$. Hence, naturally the following question appears.

Question 3.5. *Let X and Z be topological spaces such that the diagonal of every mapping $f \in CC(X \times X, Z)$ is of the first Baire class. Does the inclusion $CC(X \times X, Z) \subseteq B_1(X^2, Z)$ hold?*

4. Necessary conditions on diagonals of CL-mappings

Let X and Z be metric spaces. A mapping $f : X \to Z$ is called *Lipschitz* if the Lipschitz constant

$$\mathrm{Lip}(f) = \sup_{x,y\in X, x\neq y} \frac{|f(x) - f(y)|_Z}{|x - y|_X}$$

is finite. We say that f *is Lipschitz with a constant* C if $C \geq \mathrm{Lip}(f)$. A mapping $f : X \to Z$ is *Lipschitz on a set* $A \subseteq X$ *(with a constant* $C \geq 0$*)* if $f|_A$ is Lipschitz (with a constant C). The set of all Lipschitz mappings $f : X \to Z$ we denote by $L(X, Z)$.

A mapping $f : X \to Z$ is called *Lipschitz at a point* $x \in X$ if the pointwise Lipschitz constant

$$\mathrm{Lip}_x(f) = \sup_{y\in X, y\neq x} \frac{|f(x) - f(y)|_Z}{|x - y|_X}$$

is finite. We say that f *is pointsise Lipschitz* if f is Lipschitz at every point $x \in X$.

Proposition 4.1. *Let X be a topological space, Y, Z metric spaces, $g : X \to Y$ a continuous mapping, $C, \delta > 0$ and $f : X \times Y \to Z$ a mapping which is continuous with respect to the first variable and such that for any $x \in X$ and $y \in Y$ with $|y - g(x)|_Y < \delta$ the inequality*

$$|f(x, y) - f(x, g(x))|_Z \leq C|y - g(x)|_Y$$

holds. Then the mapping $h : X \to Z$, $h(x) = f(x, g(x))$, is continuous.

Proof. Fix $x_0 \in X$ and $\varepsilon > 0$. Denote $y_0 = g(x_0)$ and $z_0 = f(x_0, y_0) = h(x_0)$. Using the continuity of the mappings g and f_{y_0} at the point x_0 we choose a neighborhood U of x_0 such that $|g(x) - y_0|_Y < \min\{\frac{\varepsilon}{2C}, \delta\}$ and $|f(x, y_0) - z_0|_Z < \frac{\varepsilon}{2}$ for every $x \in U$. Now for every $x \in U$, using the hypothesis of the proposition, we conclude that

$$|h(x) - z_0)|_Z = |f(x, g(x)) - z_0|_Z \leq |f(x, g(x)) - f(x, y_0)|_Z + |f(x, y_0) - z_0|_Z <$$
$$< C|g(x) - y_0|_Y + \frac{\varepsilon}{2} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

Hence, h is continuous at the point x_0. □

Let X and Y be topological spaces. A mapping $f : X \to Y$ is said to be *σ-continuous* (see [4]), if there exists a sequence $(A_n)_{n=1}^{\infty}$ of closed sets $A_n \subseteq X$ such that all the restrictions $f|_{A_n}$ are continuous. In [3] such functions are called *piecewise continuous*.

The following result for CL-mappings on a product of a Baire and a metric spaces was, in fact, established in [7, Theorem 2].

Theorem 4.2. *Let X be a topological space, Y, Z metric spaces, $g : X \to Y$ a continuous mapping and $f : X \times Y \to Z$ a mapping which is continuous with respect to the first variable and pointwise Lipschitz with respect to the second variable at every point of the set $E = \{(x, g(x)) : x \in X\}$. The the mapping $h : X \to Z$, $h(x) = f(x, g(x))$, is σ-continuous.*

Proof. For every $n \in \mathbb{N}$ we denote by A_n the set of all points $x \in X$ such that for every $y \in Y$ with $|y-g(x)|_Y < \frac{1}{n}$ the inequality

$$|f(x, y) - f(x, g(x))|_Z \leq n|y - g(x)|_Y$$

holds. We show that $\overline{A_n} \subseteq A_{2n}$ for every $n \in \mathbb{N}$.

To obtain a contradiction, suppose that $x_0 \notin A_{2n}$. We show that $x_0 \notin \overline{A_n}$. We choose $y \in Y$ such that $|y - y_0|_Y < \frac{1}{2n}$ and $|f(x_0, y) - f(x_0, y_0)|_Z > 2n|y - y_0|_Y$, where $y_0 = g(x_0)$. The continuity of the mapping $\varphi(x) = \frac{|f(x,y)-f(x,y_0)|_Z}{|y-g(x)|_Y}$ at x_0 implies the existence of a neighborhood U_1 of x_0 such that $\varphi(x) > 2n$ for every $x \in U_1$. Notice that $|y-y_0|_Y > 0$. Therefore, the continuity of the mapping $\psi(x) = |g(x)-y_0|_Y-|y-g(x)|_Y$ at x_0 and the condition $\psi(x_0) < 0$ imply the existence of a neighborhood U_2 of x_0 such that $|g(x)-y_0|_Y < |y-g(x)|_Y$ and $|g(x) - y_0|_Y < \frac{1}{2n}$ for every $x \in U_2$. We set $U_0 = U_1 \cap U_2$ and prove that $U_0 \cap A_n = \emptyset$.

Let $x \in U_0$. Since $|f(x, y)-f(x, y_0)|_Z \leq |f(x, y)-f(x, g(x))|_Z+|f(x, g(x))-f(x, y_0)|_Z$, there exists $\tilde{y} \in \{y, y_0\}$ such that $|f(x, \tilde{y})-f(x, g(x))|_Z \geq \frac{1}{2}|f(x, y)-f(x, y_0)|_Z$. Remark that $|y-g(x)|_Y \leq |y-y_0|_Y+|y_0-g(x)|_Y < \frac{1}{n}$ and $|y_0 - g(x)|_Y < \frac{1}{2n} < \frac{1}{n}$. Hence, $|\tilde{y} - g(x)|_Y < \frac{1}{n}$ and

$$\frac{|f(x, \tilde{y}) - f(x, g(x))|_Z}{|\tilde{y} - g(x)|_Y} \geq \frac{1}{2}\frac{|f(x, y) - f(x, y_0)|_Z}{|y - g(x)|_Y} = \frac{1}{2}\varphi(x) > n.$$

Therefore, $U_0 \cap A_n = \emptyset$ and $\overline{A_n} \subseteq A_{2n}$.

For every $n \in \mathbb{N}$ let $F_n = \overline{A_n}$. Since $F_n \subseteq A_{2n}$, Proposition 4.1 implies the continuity of $h|_{F_n}$. It remains to observe that $\bigcup_{n=1}^{\infty} F_n = X$. □

Corollary 4.3. *Let X, Z be metric spaces and let $f : X^2 \to Z$ be a mapping which is continuous with respect to the first variable and pointwise Lipschitz with respect to the second one at every point of the diagonal $\Delta = \{(x, x) : x \in X\}$. Then the mapping $g : X \to Z$, $g(x) = f(x, x)$, is σ-continuous.*

5. Characterization of diagonals of CL-mappings

We first prove several auxiliary facts.

Lemma 5.1. *Let X be a metric space and G an open set in X^2 such that $\Delta = \{(x, x) : x \in X\} \subseteq G$. Then there exists a continuous function $g : X \to (0, +\infty)$ such that $\{(x, y) \in X^2 : |x - y|_X < g(x)\} \subseteq G$.*

Proof. For every point $x \in X$ we choose a neighborhood U_x of x in X and a number $\varepsilon_x > 0$ such that for any $y \in U_x$ the inclusion

$$\{(x', y') \in X^2 : |x' - y|_X < \varepsilon_x, |y' - y|_X < \varepsilon_x\} \subseteq G$$

holds. Let $(V_i : i \in I)$ be a locally finite open cover of X which refines $(U_x : x \in X)$, and $(\varphi_i : i \in I)$ a locally finite partition of unity on X subordinated to $(V_i : i \in I)$. For every $i \in I$ we choose $x \in X$ such that $V_i \subseteq U_x$ and let $\delta_i = \varepsilon_x$. It is easy to verify that the function $g : X \to (0, +\infty)$, $g(x) = \sum_{i \in I} \delta_i \varphi_i(x)$, is to be found. □

Lemma 5.2. *Let X be a metric space, $(H_n)_{n=1}^{\infty}$ a sequence of open in X^2 sets H_n such that $\Delta = \{(x, x) : x \in X\} \subseteq H_n$ for every $n \in \mathbb{N}$. Then there exist a sequences $(G_n)_{n=1}^{\infty}$ and $(F_n)_{n=1}^{\infty}$ of open sets G_n and closed sets F_n in X^2 and sequences $(\varphi_n)_{n=1}^{\infty}$ and $(\delta_n)_{n=1}^{\infty}$ of separately Lipschitz functions $\varphi_n : X^2 \to [0, 1]$ and positive functions $\delta_n : X \to (0, +\infty)$ with the following conditions:*

1. *$\Delta \subseteq G_{n+1} \subseteq F_n \subseteq G_n \subseteq H_n$ for every $n \in \mathbb{N}$;*
2. *$X^2 \setminus G_n \subseteq \varphi_n^{-1}(0)$ and $F_n \subseteq \varphi_n^{-1}(1)$ for every $n \in \mathbb{N}$;*
3. *$|u - v|_X \geq \delta_n(x)$ for every $x \in X$, $n \in \mathbb{N}$ and $u, v \in X$ with $(x, u) \in G_{n+1}$ and $(x, v) \in X^2 \setminus F_n$.*

Proof. Using Lemma 5.1 we construct a sequence $(g_n)_{n=1}^{\infty}$ of continuous functions $g_n : X \to (0, +\infty)$ such that $g_{n+1}(x) < \frac{1}{16} g_n(x)$ for all $n \in \mathbb{N}$ and $x \in X$, and $G_n = \{(x, y) \in X^2 : |x - y|_X < g_n(x)\} \subseteq H_n$ for every $n \in \mathbb{N}$. Clearly, the sequences $(G_n)_{n=1}^{\infty}$ and $(F_n)_{n=1}^{\infty}$ of sets $F_n = \{(x, y) \in X^2 : |x - y|_X \leq \frac{1}{8} g_n(x)\}$ satisfy condition (1). Moreover the sequence $(\delta_n)_{n=1}^{\infty}$ of function $\delta_n(x) = \frac{1}{16} g_n(x)$ satisfies condition (3).

Denote by $|\cdot - \cdot|_Z$ the metric on $Z = X^2$ defined by the formula: $|z_1 - z_2|_Z = \max\{|x_1 - x_2|_X, |y_1 - y_2|_X\}$, where $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$. For any $n \in \mathbb{N}$ and $z \in Z$ let $\alpha(z) = |z - F_n|_Z$, $\beta(z) = |z - Z \setminus G_n|_Z$ and $\varphi_n(z) = \frac{\beta(z)}{\alpha(z) + \beta(z)}$. Evidently, all the functions φ_n satisfy (2). It remains to show that all the functions φ_n are separately Lipschitz.

Fix $n \in \mathbb{N}$ and $x_0 \in X$. Using the continuity of g_n at x_0, we choose a positive number $\varepsilon \leq \frac{g_n(x_0)}{8}$ such that $g_n(x) > \frac{7 g_n(x_0)}{8}$ and $g_n(x) < 2 g_n(x_0)$ whenever $|x - x_0|_X < \varepsilon$. Prove that $\max\{\alpha(z), \beta(z)\} \geq \varepsilon$ for every $z = (x_0, y) \in Z$. Suppose that there exist $y \in X$, $z_1 = (x_1, y_1) \in F_n$ and $z_2 = (x_2, y_2) \in Z \setminus G_n$ such that $|z - z_1|_Z < \varepsilon$ and $|z - z_2|_Z < \varepsilon$, where $z = (x_0, y)$. Since $|x_0 - x_1|_X < \varepsilon$ and $(x_1, y_1) \in F_n$, $|x_1 - y_1|_X \leq \frac{1}{8} g_n(x_1) < \frac{1}{4} g_n(x_0)$. Analogously, since $|x_0 - x_2|_X < \varepsilon$ and $(x_2, y_2) \in Z \setminus G_n$, $|x_2 - y_2|_X \geq g_n(x_2) > \frac{7 g_n(x_0)}{8}$. Now

$$|y_2 - y_1|_X \geq |y_2 - x_2|_X - |x_2 - x_0|_X - |x_0 - x_1|_X - |x_1 - y_1|_X \geq \frac{3}{8} g_n(x_0).$$

But

$$|y_2 - y_1|_X \leq |y_2 - y|_X + |y - y_1|_X \leq \frac{1}{4} g_n(x_0),$$

a contradiction.

Hence, $\max\{\alpha(z), \beta(z)\} \geq \varepsilon$ for every $(x_0, y) \in Z$. Since the functions α, β and the function $\varphi : (0, +\infty)^2 \setminus [0, \varepsilon]^2 \to \mathbb{R}$, $\varphi(s, t) = \frac{t}{s+t}$, are Lipschitz and $\varphi_n(z) = \varphi(\alpha(z), \beta(z))$ for every $z = (x_0, y)$, the function φ_n is Lipschitz with respect to the first variable.

Similarly one can prove that φ_n is Lipschitz with respect to the second variable. □

We say that $f : X \to Z$ *is bounded* if there exists $C \geq 0$ such that $|f(x) - f(y)|_Z \leq C$ for every $x, y \in X$.

Proposition 5.3. *Let X and Z be metric spaces, let $f : X \to Z$ be a bounded mapping and let $A, B \subseteq X$ and $\delta > 0$ be such that $X = A \cup B$, f is Lipschitz on A and on B and $|x - y|_X \geq \delta$ for every $x \in X \setminus A$ and $y \in X \setminus B$. Then f is Lipschitz.*

Proof. Choose $C \geq 0$ such that $|f(y) - f(x)|_Z \leq C$ for every $x, y \in X$. It is easy to see that $\mathrm{Lip}(f) \leq \max\{\mathrm{Lip}(f|_A), \mathrm{Lip}(f|_B), \frac{C}{\delta}\}$. □

A topological space Z is called *an absolute extensor for a topological space* X, if for any closed set $A \subseteq X$ every continuous mapping $g : A \to Z$ can be extended to a continuous mapping $f : X \to Z$.

Let X and Z be topological spaces. A mapping $f : X \to Z$ is said to be *a mapping of the stable first Baire class*, if there exists a sequence $(f_n)_{n=1}^{\infty}$ of continuous mappings $f_n : X \to Z$ which pointwise stably converges to f, i.e. for every $x \in X$ the sequence $(f_n(x))_{n=1}^{\infty}$ stably converges to $f(x)$. The collection of all mappings $f : X \to Z$ of the stable first Baire class we denote by $B_1^d(X, Z)$.

Theorem 5.4. *Let X be a metric space, (Z, λ) a metric equiconnected space, where λ is Lipschitz with respect to third variable and $g \in B_1^d(X, Z)$. Then there exists a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and Lipschitz with respect to the second one.*

Proof. Let $(g_n)_{n=1}^{\infty}$ be a sequence of continuous mappings $g_n : X \to Z$ which pointwise stably converges to g. Let $G_0 = F_0 = X^2$ and

$$H_n = \{(x, y) \in X^2 : |g_n(x) - g_n(y)|_Z < \frac{1}{n}, |g_{n+1}(x) - g_{n+1}(y)|_Z < \frac{1}{n}\}$$

for every $n \in \mathbb{N}$. By Lemma 5.2 there exist sequences $(G_n)_{n=1}^{\infty}$ and $(F_n)_{n=1}^{\infty}$ of open sets G_n and closed sets F_n in X^2 and sequences $(\varphi_n)_{n=1}^{\infty}$ and $(\delta_n)_{n=1}^{\infty}$ of separately Lipschitz functions $\varphi_n : X^2 \to [0, 1]$ and positive functions $\delta_n : X \to (0, +\infty)$ which satisfy conditions (1), (2) and (3) of Lemma 5.2. According to Theorem 3.1 the mapping $f : X^2 \to Z$ defined by (3.1) is separately continuous. Clearly that g the diagonal of f. It remains to verify that f is Lipschitz with respect to the second variable.

Fix $x_0 \in X$ and show that the function $h = f^{x_0} : X \to Z$, $h(y) = f(x_0, y)$, is Lipschitz. It follows from (3.2) that

$$h(y) = \lambda(g_n(x_0), g_{n+1}(x_0), \varphi_n(x_0, y)) \tag{5.1}$$

for every $n \in \mathbb{N}$ and $y \in A_n = \{y \in X : (x_0, y) \in F_{n-1} \setminus G_{n+1}\}$. Since for every $n \in \mathbb{N}$ the set $h(A_n)$ contained in the compact set $\{\lambda(g_n(x_0), g_{n+1}(x_0), t) : t \in [0, 1]\}$, the mapping $h|_{A_n}$ is bounded. Using (i), we choose a number $n_0 \in \mathbb{N}$ such that $g_n(x_0) = g(x_0)$ for all $n > n_0$. Then $h(X) = \bigcup_{n=1}^{n_0} h(A_n)$ and h is bounded.

Since λ is Lipschitz with respect to the third variable and φ_n is Lipschitz with respect to the second variable, it follows from (5.1) that h is is Lipschitz on each set A_n. For every $n \in \mathbb{N}$ put $B_n = \bigcup_{k=1}^{n} A_k$. Note that For every $n \in \mathbb{N}$ if $u \in A_{n+1} \setminus B_n$ and $v \in B_n \setminus A_{n+1}$ then $(x_0, u) \in G_{n+1}$, $(x_0, v) \in X^2 \setminus F_n$ and $|u - v|_X \geq \delta_n(x_0)$ according to (3) of Lemma 5.2. Now it easy to verify by induction on n that according to Proposition 5.3 h is Lipschitz on each set B_n. Moreover $h(y) = g(x_0)$ for every $y \in A = \{u \in X : (x_0, u) \in F_{n_0}\}$, in particular, h is Lipschitz on A. Therefore by Proposition 5.3, h is Lipschitz on $X = A \cup B_{n_0}$. □

Theorem 5.5. *Let X be a metric space, (Z, λ) a metric equiconnected space, where λ is Lipschitz with respect to third variable and Z is an absolute extensor for X, and let $g : X \to Z$ be a mapping. Then the following conditions are equivalent:*

(i) $g \in B_1^d(X, Z)$;

(ii) there exists a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and Lipschitz with respect to the second one;

(iii) there exists a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and pointwise Lipschitz with respect to the second variable at every point of the diagonal $\Delta = \{(x, x) : x \in X\}$.

Proof. The implication $(i) \Rightarrow (ii)$ follows from Theorem 5.4.

The implication $(ii) \Rightarrow (iii)$ is obvious.

$(iii) \Rightarrow (i)$. By Corollary 4.3, g is σ-continuous. Now according to [3, Theorem 6.3], $g \in B_1^d(X, Z)$. □

The following questions concern the essentiality of come conditions in Theorems 5.5.

Question 5.6. *Let X be a metric space, (Z, λ) a metric equiconnected space and $f : X^2 \to Z$ a mapping which is continuous with respect to the first variable and Lipschitz with respect to the second one. Does the mapping $g : X \to Z$, $g(x) = f(x, x)$, belong to the first stable Baire class?*

6. Characterization of diagonals of CD-mappings

Let X and Y be normed spaces. A mapping $f : X \to Y$ is *Fréchet differentiable at a point* $x_0 \in X$ if there exists a linear continuous operator $T : X \to Y$ such that

$$\lim_{t\to 0} \frac{f(x_0 + tx) - f(x_0) - T(x)}{t} = 0.$$

If $f : X \to Y$ is Fréchet differentiable at every point $x \in X$, then f is called *Fréchet differentiable*.

Lemma 6.1. *Let X be a normed space with (continuously) Fréchet differentiable norm, $(H_n)_{n=1}^{\infty}$ a sequence of open in X^2 sets H_n such that $\Delta = \{(x, x) : x \in X\} \subseteq H_n$ for every $n \in \mathbb{N}$. Then there exist a sequence $(G_n)_{n=1}^{\infty}$ of open in X^2 sets G_n and a sequence $(\varphi_n)_{n=1}^{\infty}$ of jointly continuous, Lipschitz and (continuously) differentiable with respect to the second variable functions $\varphi_n : X^2 \to [0, 1]$ with the following conditions:*

(1) $\Delta \subseteq G_{n+1} \subseteq \overline{G_{n+1}} \subseteq G_n \subseteq H_n$ *for every* $n \in \mathbb{N}$;
(2) $X^2 \setminus G_n \subseteq \varphi_n^{-1}(0)$ *and* $\overline{G_{n+1}} \subseteq \varphi_n^{-1}(1)$ *for every* $n \in \mathbb{N}$.

Proof. We first using Lemma 5.1 construct a sequence $(g_n)_{n=1}^{\infty}$ of continuous functions $g_n : X \to (0, +\infty)$ such that $g_{n+1}(x) < g_n(x)$ for all $n \in \mathbb{N}$ and $x \in X$, and $G_n = \{(x, y) \in X^2 : \|x - y\| < g_n(x)\} \subseteq H_n$ for every $n \in \mathbb{N}$, and choose a continuously differentiable function $\psi : \mathbb{R} \to [0, 1]$ such that $\psi(x) = 1$ if $x \le 0$ and $\psi(x) = 0$ if $x \ge 1$. It remains to observe that $\varphi_n(x, y) = \psi\left(\frac{\|x-y\|-g_{n+1}(x)}{g_n(x)-g_{n+1}(x)}\right)$ for all $n \in \mathbb{N}$ and $x, y \in X$. □

Theorem 6.2. *Let X be a normed space with a (continuously) Fréchet differentiable norm, Z a normed space and $g : X \to Z$. Then the following conditions are equivalent:*

(i) $g \in B_1^d(X, Z)$;

(ii) there exists a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and is Lipschitz and continuously Fréchet differentiable with respect to the second variable;

(iii) there exists a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and pointwise Lipschitz with respect to the second variable at every point of the diagonal $\Delta = \{(x, x) : x \in X\}$.

Proof. *(i)* $\Rightarrow$ *(ii)*. Let $(g_n)_{n=1}^{\infty}$ be a sequence of continuous mappings $g_n : X \to Z$ which pointwise stably converges to g. Let $G_0 = F_0 = X^2$ and

$$H_n = \{(x, y) \in X^2 : |g_n(x) - g_n(y)|_Z < \frac{1}{n},\ |g_{n+1}(x) - g_{n+1}(y)|_Z < \frac{1}{n}\}$$

for every $n \in \mathbb{N}$. By Lemma 6.1 there exist a sequence $(G_n)_{n=1}^{\infty}$ of open on X^2 sets G_n and a sequence $(\varphi_n)_{n=1}^{\infty}$ of jointly continuous, Lipschitz and (continuously) differentiable with respect to the second variable functions $\varphi_n : X^2 \to [0, 1]$ which satisfy conditions (1) and (2) of Lemma 6.1. For every $n \in \mathbb{N}$ let $F_n = \overline{G_{n+1}}$. According to Theorem 3.1 the mapping $f : X^2 \to Z$ defined by (3.1), where $\lambda(x, y, t) = (1 - t)x + ty$, is separately continuous.

Remark that $\varphi_n(x, y)\varphi_{n+1}(x, y) = \varphi_{n+1}(x, y)$ for every $n \in \mathbb{N}$, therefore for the normed space Y condition (3.2) is of the following form:

$$f(x, y) = (1 - \varphi_n(x, y))g_n(x) + \varphi_n(x, y)g_{n+1}(x) -$$

$$-\varphi_{n+1}(x, y)g_{n+1}(x) + \varphi_{n+1}(x, y)g_{n+2}(x) \tag{6.1}$$

for all $(x, y) \in F_{n-1} \setminus F_{n+1}$.

Now it follows from the properties of functions φ_n that f is (continuously) Fréchet differentiable and Lipschitz with respect to the second variable on the open set $G_0 \setminus F_1 = F_0 \setminus F_1$ and on every open set $G_n \setminus F_{n+1}$, $n \in \mathbb{N}$. Hence, f is (continuously) Fréchet differentiable and Lipschitz with respect to the second variable on the open set $X^2 \setminus E$, where $E = \bigcap_{n=1}^{\infty} G_n$ (the and Lipschitz property can be obtained by Proposition 5.3).

Fix $(x_0, y_0) \in E$ and choose a number $n_0 \in \mathbb{N}$ such that $g_n(x_0) = g(x_0)$ for all $n > n_0$. Then $f(x_0, y) = g(x_0)$ for al $y \in \{x \in X : (x_0, x) \in G_{n_0+1}\}$. Therefore, f is (continuously) Fréchet differentiable with respect to the second variable at (x_0, y_0). Moreover f^{x_0} is Lipschitz by Proposition 5.3.

The implication *(ii)* $\Rightarrow$ *(iii)* is obvious.

(iii) $\Rightarrow$ *(i)*. By Corollary 4.3, g is σ-continuous. Since by Dugundji theorem normed space Z is an absolute extensor for X, $g \in B_1^d(X, Z)$ by [3, Theorem 6.3]. □

Question 6.3. *Let X, Z be normed spaces and $g \in B_1^d(X, Z)$. Does there exist a mapping $f : X^2 \to Z$ with the diagonal g which is continuous with respect to the first variable and is Lipschitz and continuously Fréchet differentiable with respect to the second one?*

Question 6.4. *Let X, Z be normed spaces and $f : X^2 \to Z$ a mapping which is continuous with respect to the first variable and (weakly) Gâteaux differentiable with respect to the second one. Does the mapping $g : X \to Z$, $g(x) = f(x, x)$, belong to the first stable Baire class?*

References

[1] Baire R. *Sur les fonctions de variables reélles*, Ann. Mat. Pura Appl., **3** (3) (1899), 1–123.

[2] Banakh T. *(Metrically) Quater-stratifable spaces and their applications in the theory of separately continuous functions*, Mat. Stud. **18** (1) (2002), 10–18.

[3] Banakh T., Bokalo B. *On scatteredly continuous maps between topological spaces*, Top. Appl. **157** (1) (2010), 108–122.

[4] Banakh T., Kutsak S., Maslyuchenko O., Maslyuchenko V. *Direct and inverse problems of the Baire classification of integrals dependent on a parameter*, Ukr. Math. J., **56**:11 (2004), 1443–1457 (in Ukrainian).

[5] Burke M. *Borel measurability of separately continuous functions*, Top. Appl. **129** (1) (2003), 29–65.

[6] Hahn H. *Theorie der reellen Funktionen 1*, Berlin: Verlag von Julius Springer (1921).

[7] Herasymchuk V., Maslyuchenko V., Mykhaylyuk V. *Varieties of Lipschitz condition and discontinuity point sets of separately differentiable functions*, Bull. of Chernivtsi National University. Mathematics, **134** (2002), 22–29 (in Ukrainian).

[8] Karlova O. *Separately continuous σ-discrete mappings*, Bull. of Chernivtsi National University. Mathematics, **314-315** (2006), 77–79 (in Ukrainian).

[9] Lebesque H. *Sur l'approximation des fonctions*, Bull. Sci. Math. **22** (1898), 278–287.

[10] Lebesque H. *Sur les fonctions respresentables analytiquement*, Journ. de Math. **2** (1) (1905), 139–216.

[11] Maslyuchenko V., Mykhaylyuk V., Sobchuk O. *Construction of a separately continuous function of n variables with the given diagonal*, Math. Stud. **12** (1) (1999), 101–107 (in Ukrainian).

[12] Maslyuchenko O.V., Maslyuchenko V.K., Mykhaylyuk V.V., Sobchuk O.V. *Paracompactness and separately continuous mappings*, General Topology in Banach spaces, Nova Sci. Publ., Nantintong - New-York. (2001), 147–169.

[13] Moran W. *Separate continuity and support of measures*, J. London. Math. Soc. **44** (1969), 320–324.

[14] Mykhaylyuk V. *Construction of separately continuous functions of n variables with the given resriction*, Ukr. Math. Bull. **3** (3) (2006), 374–381 (in Ukrainian).

[15] Rudin W. *Lebesgue's first theorem*, Math. Analysis and Applications, Part B. Adv. in Math. Supplem. Studies, **7B** (1981), 741–747.

[16] Vera G. *Baire measurability of separately continuous functions*, Quart. J. Math. Oxford. **39**(153) (1988), 109–116.